



Variationally computed ro-vibrational energies of sulphur trioxide

Underwood, Daniel; Yurchenko, Sergei N.; Tennyson, Jonathan; Fateev, Alexander

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Underwood, D., Yurchenko, S. N., Tennyson, J., & Fateev, A. (2012). *Variationally computed ro-vibrational energies of sulphur trioxide*. Poster session presented at 22nd International Conference on High Resolution Molecular Spectroscopy, Prague, Czech Republic.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Variationally Computed Ro-Vibrational Energies of Sulphur Trioxide

Daniel Underwood, Sergei N. Yurchenko, Jonathan Tennyson, Alexander Fateev

Department of Physics and Astronomy, University College London,
Gower Street, London WC1E 6BT



Department of Chemical and Biochemical Engineering,
Technical University of Denmark, 2800 Kgs. Lyngby,
Denmark



UCL

Introduction

We present preliminary results of a collaboration between academic and industrial efforts.

We aim to create a high-temperature spectroscopic database for SO_3 which is essential in the analysis of complex data sets obtained through *in situ* measurements on industrial fuel units in power plants.



Ultimately, the results will be used in working to:

- increase the efficiency of electricity production by reducing NO_x emissions;
- optimising Ammonia consumption (Ammonia is used as a reductant);
- reducing material corrosion and;
- prolonging lifetimes of the selective catalytic reduction units.

Atmospheric & Astronomical significance:


SO_3 is also a contributor to the formation of acid rain, and plays a significant role in atmospheric chemistry; it is also believed to be a constituent of atmospheres in astronomical environments, such as Venus and Io [1].

The Molecule

- A trigonal planar molecule that belongs to the D point group.

- Has 4 fundamental vibrational modes:

- A' symmetric stretch
- A'' out-of-plane bend
- E' antisymmetric stretch
- E' in-plane bend



Quantum Numbers: J, K, v_1 , v_2 , $v_3^{l_3}$, $v_4^{l_4}$

- Is composed entirely of spin-0 Bosons and therefore has only a single nuclear spin configuration and must exhibit an overall symmetric wavefunction.

- We therefore only have allowed ro-vibrational transitions between states of A' and A'' symmetries.

- SO_3 is a relatively heavy molecule with a small rotational constant ($B = 0.34854333(5)\text{cm}^{-1}$ [2]). Calculations of ro-vibrational energies up to high values of J are therefore necessary for large coverage in the production of synthetic spectra.

Computational Background and TROVE

- Potential Energy Surface used in this work computed using MOLPRO.

- Re-adjusted to an experimentally determined equilibrium bond length (1.41732 Angstroms [3]).

- PES is then used by TROVE [4] to calculate ro-vibrational wavefunctions and energy calculations.

- For SO_3 we define an active space such to maximise computational efficiency by the truncation of the Hamiltonian matrix blocks. This is based on the following:

- Calculations of ro-vibrational energies up to and including $10,000\text{cm}^{-1}$;
- SO_3 exhibits a polyad structure, and thus we can employ the polyad number [2]. We impose the condition on the quantum numbers that $P = 2(v_1 + v_3) + v_2 + v_4 \leq 14$
- Computational time is further reduced by noting that only energy levels of A' and A'' symmetry are required.

Results


Fundamental Vibrations & Some Overtone/Combinations

	Experimental [2]	This Work	Martin [3]
v1	1064.924 cm^{-1}	1065.738 cm^{-1}	1064.219 cm^{-1}
v2	497.567 cm^{-1}	498.475 cm^{-1}	487.100 cm^{-1}
v3	1391.520 cm^{-1}	1387.428 cm^{-1}	1386.854 cm^{-1}
v4	530.086 cm^{-1}	528.580 cm^{-1}	527.321 cm^{-1}
2v2	995.018 cm^{-1}	995.346 cm^{-1}	970.366 cm^{-1}
2v3(l=2)	2777.871 cm^{-1}	2769.954 cm^{-1}	2770.080 cm^{-1}
2v4(l=0)	1059.814 cm^{-1}	1056.440 cm^{-1}	1054.702 cm^{-1}
2v4(l=2)	1060.452 cm^{-1}	1057.332 cm^{-1}	1055.383 cm^{-1}
v2 + v4	1027.902 cm^{-1}	1027.325 cm^{-1}	1013.453 cm^{-1}

Pure Rotational Energies

J	K	Experimental [2]	This Work	Difference
2	0	2.091 cm^{-1}	2.091 cm^{-1}	0.0003 cm^{-1}
4	0	6.970 cm^{-1}	6.971 cm^{-1}	0.001 cm^{-1}
6	0	14.638 cm^{-1}	14.640 cm^{-1}	0.002 cm^{-1}
8	0	25.093 cm^{-1}	25.097 cm^{-1}	0.003 cm^{-1}
10	0	38.336 cm^{-1}	38.341 cm^{-1}	0.005 cm^{-1}
20	0	146.333 cm^{-1}	146.354 cm^{-1}	0.021 cm^{-1}
40	0	323.876 cm^{-1}	323.923 cm^{-1}	0.046 cm^{-1}
60	0	1271.537 cm^{-1}	1271.704 cm^{-1}	0.167 cm^{-1}
70	30	1569.767 cm^{-1}	1569.969 cm^{-1}	0.201 cm^{-1}

Comparison of Reduced Energies ($E_{jk}' - E_{jk}''$) in 1550-1610 cm^{-1} Region



Next Step...

- We aim to complete the current work by calculating all ro-vibrational energies up to J = 100.

- To spectroscopically refine a new PES computed at the aug-cc-pVQZ-f12 level of theory.

- Calculations of an *ab initio* Dipole Moment Surface have already begun, using a aug-cc-pVTZ-f12 basis set, with preliminary intensity results for transitions up to J = 20. We will eventually calculate intensities up to J = 100 (T = 300K).

References

- 1) E. T.H. Chrysostom, N. Vulpanovici, T. Masiello, J. Barber, J. W. Nibler, A. Weber, A. Maki, T. A. Blake, "Coherent Raman and Infrared Studies of Sulphur Trioxide", *J. Mol. Spec.*, **210**, 233-239 (2001)
- 2) A. Maki, T. A. Blake, R. L. Sams, J. Frieh, J. Barber, T. Masiello, E. T.H. Chrysostom, J. W. Nibler, A. Weber, "Analysis of some combination-overtone infrared bands of $^{32}\text{S}^{16}\text{O}_3$ ", *J. Mol. Spec.*, **225**, 109-122 (2004)
- 3) Martin, J. M. L., "A fully ab initio quartic force field of spectroscopic quality for SO_3 ", *Spectrochimica Acta Part A*, **55**, 105-112 (2002)
- 4) S. N. Yurchenko, W. Thiel, P. Jensen, "Theoretical ROVibrational Energies (TROVE): A robust numerical Approach to the calculation of rovibrational energies for polyatomic molecules", *J. Mol. Spec.*, **245**, 126-140 (2007)

Acknowledgements

The projects are supported by the ERC grant (Advanced Investigator Project 267219) and by Energinet.dk (Denmark, Project 10442).